1st Midsemester Test 2020. 04. 06.

B

traveling at 40.0	m/s just before it st	trikes a 40.0-g golf	ball at rest on a te	lub of mass 250 g is e. After the collision, l of the golf ball just
a. 20,3 m/s	b. 80,2 m/s	c. 62,5 m/s	d. 56,7 m/s	e. none of them
	s from rest and rota s in 3.00 s. The ang		_	n to reach an angular
a. $4 \frac{1}{s^2}$	b. 4 1/s ²	c. 2,5 1/s ²	d. 6 1/s ²	e. none of them
rad/s in the begi		g wheel rotates 2.0	00 s and finally its	ngular velocity is 20 angular speed is 64
a. 60 rad	b. 84 rad	c. 50 rad	d. 40 rad	e. none of them
	from rest and rotat l/s in 4.00 s. Find th		_	to reach an angular states in this time:
a. 20 rad	b. 4 rad	c. 32 rad	d. 18 rad	e. none of them
	tal force of 200.0			tarted from rest by a of the disk. Find the
a. 2,5 1/s ²	b. 6,2 1/s ²	c. 1,25 1/s ²	d. 0,625 1/s ²	e. none of them
	id disc of mass 20. of mass has a speed	_		contal surface. At the rgy of the disk!
a. 750 J	b. 1500 J	c. 3000 J	d. 375 J	e. none of them
m, and the object	et is released from	rest there. It proce	eds to move without	ly stretched by 0.200 out friction. The next speed of the object?
a. 0,253 m/s	b. 3,14 m/s	c. 0,628 m/s	d. 1,356 m/s	e. none of them
	rticle moves in the entum of the partic			$+6\vec{j}$ m/s. Determine $=5\vec{i}$:
a. 45 k	b. 30 k	c. 15 i	d. 6 i	e. none of them

9. A 2000-kg car stopped at a traffic light is struck from the rear by a 1200-kg car, and the two become entangled, moving along the same path as that of the originally moving car. If the smaller car were moving at 40.0 m/s before the collision, what is the velocity of the entangled cars after the collision?							
a. 22 m/s	b. 15 m/s	c. 16 m/s	d. 6,67 m/s	e. none of them			
10. A horizontal 600-N merry-go-round is a solid disk of radius 2.0 m, and it rotates at 4 rad/s. Find the kinetic energy of the disk:							
a. 960 J	b. 2560 J	c. 3000 J	d. 1280 J	e. none of them			
11. The angular position of the particle is $\Theta(t)$ = 2 - 3 t^2 + 2 t^3 (rad), its average angular velocity between 1s and 3s is:							
a. 22 1/s	b. 23 1/s	c. 16 1/s	d. 25 1/s	e. none of them			
12. A ball of mass 0.15 kg is released from a height of 1.25 m. It rebounds from the floor to reach a height of 0.80 m. What impulse was given to the ball by the floor?							
a. 1,35 Ns	b. 0,6 Ns	c. 0,75 Ns	d. 1,82 Ns	e. none of them			
13. The angular velocity of the particle is $\omega(t)=2+3t^2+t^3$ (1/s), its average angular acceleration between 1s and 3s is:							
a. $50 \ 1/s^2$	b. 28 1/s ²	c. 6 1/s ²	d. 25 1/s ²	e. none of them			
14. The simple pendulum has a string of 0,25 m. The period of the harmonic motion of the pendulum is:							
a. 1,0 s	b. 1,4 s	c. 2,9 s	d. 0,69 a	e. none of them			
15. A 4.00 kg particle has a velocity of $-6\vec{i} + 8\vec{j}$ m/s. The magnitude of the linear momentum:							
a. 24 Ns	b. 15 Ns	c. 40 Ns	d. 32 Ns	e. none of them			
16. A rotating wheel requires 4.00 s to rotate through 320 rad. Its angular speed at the end of the 4.00 s interval is 40 rad/s. What is the magnitude of the constant angular acceleration of the wheel?							
a. 10 1/s ²	b. 120 1/s ²	c. 20 1/s ²	d. 160 1/s ²	e. none of them			